Foot and Mouth Disease in Cameroon: A systematic review to support its progressive control

*1Lendzele SS, ²Aubin KA and ²Mavoungou JF

¹Organisation for Milk and Meat Production (PLEB), Adamawa, Cameroon ²Department of Biology and Animal Ecology, Institute of Research in Tropical Ecology, Libreville, Gabon

Corresponding author: sevidzem.lendze@gmail.com

Received on: 31/01/2022	Accepted on: 11/05/2022	Published on: 17/05/2022
-------------------------	-------------------------	--------------------------

ABSTRACT

Foot and mouth disease (FMD) is an infectious viral transboundary disease of even-toed animals, caused by the foot and mouth disease virus (FMDV). It affects the development of the livestock sector in sub-Saharan Africa (SSA) and is endemic in Cameroon. The main objective of the study was to provide reviewed data on the epidemiology, risks, control and the existing gaps on FMD in Cameroon from 1990 to 2021 to support its progressive control. To attain this objective, relevant and available documents (n=45) on FMD in Cameroon such as peer-reviewed papers (n=27), dissertations (n=11), conferences (n=5) and technical reports (n=2) were used. Studies on the molecular and serological epidemiology, the role of carrier/mobile animals, the evaluation of immune-cross protection between serotypes and the quantification of serotype-specific transmission parameters using mathematical models have been discussed. The environmental drivers of FMD in cattle markets, abattoir and herd-level, as well as risk factors such as husbandry practices, communal grazing, regional/international livestock trading system, transhumance and fomites were presented. As international borders remain porous with poor FMD monitoring infrastructure at veterinary checkpoints in a country with no candidate vaccine, vaccination program and commercial vaccines available, farmers are left with no option other than to manage the disease with traditional formulations and veterinary pharmaceuticals mostly antibiotics. Tri-Solfen® (TS), a wound dressing formulation was efficient in managing FMD in Cameroon. The current gaps in FMD knowledge in Cameroon were presented. There was a need for a comprehensive epidemiological study in the major cattle rearing regions of Cameroon. A regional study to identify all the risk factors for the transmission of FMD was required and a socio-economic impact study of FMD was required in Cameroon.

Keywords: Foot-and-mouth disease, serotypes, epidemiology, risk factors, transmission, control, gaps analysis

Cite This Article as: Lendzele SS, Aubin KA and Mavoungou JF (2022). Foot and Mouth Disease in Cameroon: A systematic review to support its progressive control. J. Vet. Res. Adv., 04(01): 15-26.

Introduction

Foot-and-mouth disease is a highly contagious viral disease of even-toed animals belonging to the family *Picorna viridae* and genus *Aphtho virus*. It is among the diseases that greatly affect the progress of the livestock industry in SSA because its endemicity does not only reduces animal production and health but also hinders international trade with FMD free regions (James and Rushton, 2002). This disease is caused by seven known serotypes (O, A, C, Asia 1, South African Territories (SAT) 1, SAT 2, SAT 3) with characteristic ecologically distinct evolutionary lineages or subtypes within serotypes and limited cross-immunity between serotypes or lineages (Knowles and Samuel, 2003).

Cameroon is found in FMD pool 5 with four serological types (O, A, SAT 1, SAT 2) endemic to countries in the Central and West African subregions. In Cameroon, molecular studies have confirmed that serotypes-O, A, and SAT 2 are endemic (Bronsvoort et al., 2004; Bronsvoort et al., 2006; Bertram et al., 2018a). In 2019, SAT1 topotype X was reported for the first time in bovine epithelial tissues from North Cameroon (Bertram et al., 2019; Ehizibolo et al., 2020). Indeed, serological epidemiological studies have established the occurrence of five serotypes-O, A, SAT1, SAT 2, SAT 3 (Ludi et al., 2016 ; Sevidzem et al., 2019a). As international borders remain permeable without strict veterinary checkpoints to identify, isolate and prevent sick animals from entering the country, the risk of incursions of FMD is very obvious shown viruses as in studiesconducted at border levels between Cameroon and neighbouring countries such as Chad, Central African Republic (CAR), and

Copyright: Lendzele et al. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Nigeria (Ehizibolo et al., 2020; Mebanga et al., 2020; Schnell et al., 2019; Bertram et al., 2018b. Xiao et al., 2015; Lendzele et al., 2021).

The risk-based control approach using the global progressive control pathway for FMD control (PCP-FMD) is the target for Cameroon as it is currently in the PCP-FMD stage one (Ehizibolo et al., 2020).FMD transmission risk sources such as communal grazing, mobile livestock especially cattle, the role of carrier animals, role of fomites (a fomite is any material or object that can harbour an infectious disease and probably contaminate the environment and hosts), husbandry system, high livestock density, and the environment around herds, abattoirs and cattle markets have been examined and identified to be potential transmission areas to target for its mitigation (Lendzele et al., 2021; Lendzele et al., 2020 ; Sevidzem et al., 2019b; Bronsvoort et al., 2016; Motta et al., 2018).

The control of FMD in Cameroon is a major concern as no vaccination programs are ongoing and its management is mostly conducted by farmers themselves using veterinary pharmaceuticals mostly purchased from black markets andonly a few of them buy from recognised pharmacies (Sevidzem et al, 2019b). The field trial of the trivalent FMD vaccine (O, A, SAT 2) reduced clinical FMD but did not stop persistent infections (Bertram et al., 2018a). The field trial of TS, a topical formulation improved the overall demeanour of animals, hastened recovery time, wound healing time and quick return to pasture compared to animals treated with antibiotics and non-treated counterparts (Lendzele et al., 2020). The management of the disease with antibiotics could have public health consequences such as the risk of the development of antibiotic resistance (Vougat Ngom et al., 2020). Therefore, vaccination or the use of an alternative such as TS could be suggested for FMD control in Cameroon.

The present study aims at conducting a systematic review of relevant documents available from 1990 to 2021on FMD epidemiology, risk-based studies, control options and gap analysis to support its progressive control.

Materials and Methods

Study area: Cameroon is a bilingual country and is located at the northeastern end of the Gulf of Guinea between the equator and the tropics.

Cameroon is in the form of a triangle with a 700 km base and a 1200 km side with a total surface area of 475000 Km². It is found between longitude 9° to 16° east and latitude 2° to 13° north. It forms part of Central African countries, where it shares a boundary to the South with Equatorial Guinea, Gabon and Congo, to the West with Nigeria, to the East with the Central African Republic and Chad. Cameroon is a major cattle rearing region and supplier of livestock products to neighbouring countries in the Central African subregion. The major livestock rearing regions of Cameroon are the Far North, Adamawa, North and North West. The livestock population in Cameroon includes 31 million poultry, 6 million cattle, 7 million small ruminants, one million pigs, 150,000 donkeys and 15,000 horses (FAO, 2015). The distribution of cattle throughout the national territory is as such: 37.5% in the Far North, 33.9% in Adamawa, 11.6% in the North, 8% in the North West, 6.3% in the East and 2.7% in the West (Moumini, 2011). The present study focused on reviewing available data on FMD in all the regions of Cameroon.

Literature search methods and selection criteria: A webbased systematic search to get information from the published research literature (peer-reviewed papers and dissertations), the consultation of technical reports of the LANAVET-MINEPIA and conference presentations made at the European Commission for Foot-and-mouth disease Control (EuFMD) and Global Foot-and-mouth disease Research Alliance (GFRA) meetings as well as the EuFMD/FAO/OIE reports on FMD in Cameroon. The published documents were gotten via Google, Google Scholar, websites of local ResearchGate, and and international institutions working on FMD in Cameroon. Only those documents reporting on the epidemiology, transmission risk, and control measures of FMD from January 1990 to August 2021 were selected for this study. Based on the stated inclusion criteria, a total of 45 relevant and complete documents were retained for this systematic review, excluding documents that had no information on the study country and objectives.

Results and Discussion

Epidemiological findings on FMD: Foot and mouth disease is endemic in Cameroon where the health and productivity of about 7 million cattle, 3.5 million sheep, 4 million goats and 1.7 million pigs is threatened by it (FAO, 2015). Studies on FMD in Cameroon between 1931 and 1988 showed the circulation of serotypes O and A (Ekue et al., 1990).

The national veterinary laboratory (LANAVET) in Garoua isolated serotypes A, O, SAT 2, SAT 3 from cattle and pig samples between 1987 and 1998. Subsequently, studies in the Adamawa region from 2000 to 2006, highlighted the presence of serotypes SAT 2, O, and A (Bronsvoort et al., 2006). In 2006 within the framework of the Pan African Control of Epizootics(PACE)-Cameroon program, samples collected, part analysed in LANAVET, revealed the circulation of serotypes SAT 2, O, A, and were confirmed by theInstitute for Animal Health (IAH)-Pirbright. According to the Cameroon report (prepared by the LANAVET) from 2005 to 2011 to the International Organisation for Animal Health (OIE), it showed that in 2005 four serotypes (A, O, SAT 1, SAT 2) were detected, but samples from 2007 to 2011 were not typed. In 2011, samples were collected from FMD outbreak areas of Adamawa and North regions by the LANAVET staff and sent to Botswana Vaccine Institute (BVI) for virus isolation, phylogenetic analysis and vaccine matching. From the analysis, SAT 2 was isolated from samples collected from the North and Adamawa. In 2015, three serotypes (SAT 2, O, A) were identified in Ngaoundere (Bertram et al., 2018a). In general, four FMD serotypes (A, O, SAT 1, SAT2) have been confirmed to be circulating in Cameroon (Fig. 2). The serotypes identified in Cameroon have also been identified in neighbouring Chad (Abdel-Aziz et al., 2019). The SAT1 topotype X was reported for the first time in Cameroonian (samples from the North region) and Nigerian cattle (Bertram et al., 2019; Ehizibolo et al., 2020). The occurrence of this SAT 1 topotype X between the two countries is not surprising as they face the challenge of uncontrolled animal movement across permeable international borders that allows the contact of animals, thus facilitating the incursion of FMD viruses and spread. Similarly, the SAT 2 topotype VII identified in Chadian livestock was similar to that reported in 2015 in Cameroon (Abdel-Aziz et al., 2019), confirming that uncontrolled cattle movement in endemic SSA is a central mechanism for FMD introduction and distribution (Ularamu et al., 2020; Kerfua et al., 2018; Jori and Etter, 2016).

The role of small ruminants in FMD epidemiology is currently being considered in Cameroon. Studies conducted in the North and Adamawa regions found that subclinical small ruminants (goats and sheep) were infected with FMD and most seriously when they are reared with other animal species as well as graze around wildlife areas (Dickmu et al., 2019). In Cameroon, the mixed-species herd husbandry practice is very common with cattle kept alongside other ruminant species like goats, sheep and rarely pigs(FAO, 2015). The possibility of cross-transmission of FMD between animal species through contact has already been reported in Kenya (Chepkwony et al., 2021). Although the control of FMD by only vaccinating large ruminants without including small ruminants worked in certain settings, this has led to the negligence of the role of small ruminants in FMD epidemiological studies. However, the role of small ruminants in sustaining FMD transmission chains in an endemic setting is still debated.

Recent advances have been made in the epidemiological studies of FMD in Cameroon by the OSU and the University of Edinburgh (UoE) researchers through the use of mathematical modelling to project FMD transmission in the Far-North and Adamawa. Serological data from an FMD endemic setting (Far-North region of Cameroon) was used to probe strain-specific transmission immuno-dynamics (Pomeroy et al., 2015). They discovered that for serotypes SAT 2, SAT 3, O, a model assuming lifelong immunity fitted better for serotypes SAT 1 and O and this better fit model suggested that immunity could wane overtime. They noticed that serotype O had the highest force of infection (FOI) and longest duration of immunity, SAT1 and O displayed endemic dynamics, type A displayed epidemic dynamism and SAT 2, SAT 3 did not sustain local chains of transmission. Similarly, simulating the transmission of FMD among mobile herds in the Far-North region indicated that grazing areas observed in the field (≤5km radius) resulted in multiple epidemic peaks in a year (Kim et al., 2016). A study conducted on subclinical and mobile animals indicated the potential transmission risk by such groups (Schnell et al., 2019). In Adamawa, the role of carrier animals in FMD spread was examined and a redefinition of the carrier animal status for endemic settings was suggested (Bronsvoort et al., 2016). Herdsmen report on FMD has also been an important tool to track infected cases, but how accurate this might be has been verified in studies conducted in the Adamawa region. A study conducted to validate

herdsmen sensitivity and specificity in Adamawa concluded that although farmers could identify the disease with high sensitivity, this should be conducted in parallel with serological testing (Morgan et al., 2014).

From published technical documents and articles, the highest numberof serotypes was recorded in the Far-North region (O, A, SAT1, 2, 3) (Ludi et al., 2016). The North, Adamawa and North-West regions have the same number of serotypes (O, A, SAT 1, SAT 2) (Bronsvoort et al., 2003; Bronsvoort et al., 2006; Sevidzem et al., 2019a; Bertram et al., 2018b; Lendzele et al., 2021). The other regions such as Littoral and Center have low serotype diversity, while for the South-West and South regions there is no information on the circulating serotypes (Table 1).

FMD serotypes O and A were the most frequent serotypes in Cameroon as they were detected in almost all the years that samples were serotyped. From the nationwide FMD confirmed cases reported from 1990 to 2021, it could be deduced that the disease is widespread and endemic in Cameroon (Table 2).

Risk-based identification findings: The transmission life cycle of FMD has been studied and remains complex for endemic settings. Some aspects of the foot and mouth disease virus (FMDV) life cycle such as its persistence in the environment, the occurrence of subclinical animals (domestic and wild) and the possibility of transmission from small ruminants to cattle remain subjects for debate. The already identified potential risk sources for FMD transmission in the indigenous cattle population of Cameroon are cattle density, livestock trading practice, livestock environments, markets/abattoir husbandry systems, uncontrolled livestock movements, porous international borders and fomites. Some of these transmission sources have been fully described (Paton et al., 2018).

Livestock trading system: Livestock trade in Cameroon involves transactions in cattle markets between cattle owners who pay taxes to Government officials upon entry. It is reported that 32% of herders declare buying their cattle from conventional cattle markets (FAO, 2015). Trade animals are usually accompanied by other animals of the same herd (a herd is the smallest homogenously mixing unit of animals (Ludi et al., 2016). The unsold animals, plus those accompanying are returned to the herd. Livestock places such as cattle markets have been identified as risk areas for the transmission of infectious diseases (Motta et al., 2017). Such risk could occur if unsold animals and those that accompanied them to the market return with FMD viruses to contaminate the other animals in the herd.

The major production zones of cattle Cameroon are found in the North-West, Adamawa, Far-North and North regions. The animals bred in such major production areas are sold in other regions with major markets found in the cities of Yaounde, Douala, Bafoussam, Limbe, Bertoua. International export Ebolowa and markets for local production includeNigeria, Gabon, Congo and Equatorial Guinea. Another type of trading system is that which involves transit livestock from the Central African Republic, Chad, Sudan and Ethiopia that pass through Cameroon to Nigeria, Gabon, Congo and Equatorial Guinea (Fig. 1). Interestingly, 70% of Cameroonian herdsmen can easily identify FMD in animals in a cattle market but 8% would still buy an animal suffering from FMD because of their low prices.A survey in the Far-North region reported that about 10% of the cattle sold by pastoralists were reportedly sick at the time of the sale (Profitós et al., 2013). In Kenya, the purchase of sick animals from stock markets by small-scale dairy farmers have been reported as risk factor for FMD dissemination (Nyaguthii et al., 2019). From a national survey conducted in 2012, it was found that 31% of those who would buy an infected animal are from the North-West followed by the North (14%) and East (12%) (FAO, 2015). It is clear that despite having sound knowledge of the disease, farmers still sell FMD affected animals and this ugly practice promotes the spread of the disease.

Husbandry practices that promote FMD spread: Past surveys indicated that 92% of Cameroonian herders reported cases of FMD, but during outbreaks, only 30% separated sick animals from rest of herd, remaining 70% kept sick animals with rest of herd (FAO, 2015). A recent report on farmers management practices during FMD outbreak periods to curb the spread of disease in Adamawa showed that practice of herd isolation was very difficult based on free livestock grazing husbandry practice (Sevidzem et al., 2019b). It was noticed in the majority of the smallholder dairy farms in Adamawa that most herders used the

Table 1. FMD serotypes within the ten regions of Cameroon from 1990 to 2021.

Region	Circulating serotypes between 1931 and 2021	Kererence
		Sevidzem et al. (2019); Bertram et al. (2018a); Bronsvoort et al.
Adamawa	0, A, SAT 1, SAT 2	(2004) ; Bronsvoort et al. (2006); FAO (2015); Ekue et al. (1990)
Center	SAT 2	FAO (2015)
East	SAT 2, O	FAO (2015); Colenutt (2020)**
		Ludi et al. (2014); Pomeroy et al. (2015); Bertram et al.
Far North	O, A, SAT 1, SAT 2, SAT 3	(2018b); Fox (2017)
Littoral	0	Colenutt (2020)**
		Dickmu et al. (2019); Ehizibolo et al. (2020); Bertram et al.
North	O, A, SAT1, SAT 2	(2019)
North West	O, A, SAT 1, SAT 2	FAO (2015)
West	O, SAT 1, SAT 3	FAO (2015)
South West	Not typed*	FAO (2015)
South	Not typed*	FAO (2015)

Table 2. EMD	confirmed	construnce in	animal	anacios fre	1000 ± 2021
Table 2. FIMD	commea	serotypes m	annnais	species inc	1111990102021.

Reported year	Animal species affected	Circulating serotypes	Reference
1931 to 1959	Cattle	0	Ekue et al. (1990)
1975	Cattle	А	Ekue et al. (1990)
1976	Cattle	А	Ekue et al. (1990)
1980	Cattle	0	Ekue et al. (1990)
1981 to 1984	Cattle, Pigs	0	Ekue et al. (1990)
1985 to 1987	Cattle, Pigs, Goats, sheep	A, SAT 3	Ekue et al. (1990)
1988	Cattle, Pigs	O, A, SAT 2, SAT 3	Ekue et al. (1990)
1989	Cattle	О, А	LANAVET
1996	Cattle	O, A, SAT 2	LANAVET
1998	Cattle	SAT 2	LANAVET
2004	Cattle	О, А	Bronsvoort et al. (2004)
2005	Cattle	O, A, SAT 1, SAT 2	LANAVET
2006	Cattle	O, A, SAT 1, SAT 2, SAT 3	LANAVET; Bronsvoort et al. (2006)
2014	Cattle	O, A, SAT 1, SAT 2, SAT 3	Ludi et al. (2014)
2015	Cattle	O, A, SAT 1, SAT 2, SAT 3	Pomeroy et al. (2015); FAO (2015)
2018	Cattle	O, A, SAT 2	Bertram et al. (2018a,b)
			Sevidzem et al. (2019); Dickmu et al.
2019	Cattle, Sheep, Goats	O, A, SAT 1, SAT 2	(2019) ; Bertram et al. (2019)
2020	Cattle	SAT1	Ehizibolo et al. (2020)



Fig. 1: Livestock trade network at regional and international levels

hand-milking technique without disinfecting their hands as well as used the same utensils between animals that could promote contamination in their herds (Lendzele et al., 2021).

Despite great knowledge on FMD by Cameroonian herdsmen, the majority of them still conduct local husbandry practices that facilitate within-herd and between herds spread. Some of these practices are an exchange of bulls for reproduction purposes, allowing sick animals to mix with non-sick counterparts, transhumance towards FMD wild reservoir sites with buffaloes, antelopes, warthogs, and mixed livestock farming (whereby herders mix herds of different susceptible animal species such as goats, sheep, but rarely pigs with their cattle). Studies conducted in the North and Adamawa reported serotype O in small ruminants (sheep and goats) (Sevidzem et al., 2019a). Additionally, the study in the North identified that herders who practised grazing of small ruminants with other animal species reported high FMD contamination. Mixed livestock grazing practice leads to high animal contact and has been reported as a risk factor for FMD spread in west, central, east and southern Africa (Ekwem et al., 2021; Nyaguthii et al., 2019; Sinkala et al., 2014) and Asia (Sansamur et al., 2020). It has been highlighted that dearth of information on good management practices by herders could be attributed to the disorganized nature of the herders, where only a few of them belong to farmer organizations.

Transhumance: Transhumance is another major virus transmission mechanismand involves the southward movement of animals into game reserves, the Guinean savannah, the northwestern grasslands and the forest agro-ecological zones. These zones represent habitats for buffaloes, antelopes and warthogs. It is known that buffaloes are natural reservoir hosts for SAT 1, SAT 2, SAT 3 FMD viruses (Wekesa et al., 2015). Moreover, FMDV has already been recovered from subclinical buffaloes in South East Asia (Buckle et al., 2021). It has already been reported that transhumance herds usually come in contact with other herds at pasture, drinking and places where they camp (Schnell et al., 2019; Bronsvoort et al., 2003). In practice, transhumant animals often come in contact with other animals during the process and upon return and this could lead to contamination with new FMD viruses that are brought back to sedentary animals that did not go on transhumance. Additionally, some instances of transboundary trade herds camping overnight with native herds lead to direct contact (Pomeroy et al., 2015). Thus, transhumant animal contact could bea major transmission risk source for FMD in Cameroon.

The occurrence of wild buffalo niches in SSA permits the maintenance of FMD transmission. In some West African countries (Benin, Burkina Faso and Nigeria) and Central Africa countries (Cameroon, Central African Republic (CAR), Chad, Democratic Republic of Congo and Gabon) buffalo subspecies have been identified. For Cameroon, *Syncerus caffer brachyceros* and *Syncerus caffer nanus* occur (Di Nardo et al., 2015). Buffalo samples from Gabon and Nigeria all tested negative for FMD, but no buffalo samples were available from Cameroon for testing in this study(Di Nardo et al., 2015). However, buffalo

samples from neighbouring countries such as DR Congo, Chad and CAR were positive. Further, when wildlife species were screened in neighbouring Nigeria, FMD was detected (Atuman et al., 2020). It is well known that the Cape buffalo (*Syncerus caffer caffer*) is a long term maintenance host for SATs (Wekesa et al., 2015). Hence, crossborder transhumance where livestock share grazing, drinking and camping points with wildlife and buffaloes, exposes them to FMD infection.

The porosity of international borders : One of the FMD incursion sources is likely through transboundary trade animals from the neighbouring Central African Republic and Chad and during transit to Nigeria, Gabon and Equatorial Guinea. The importation of live animals and animal products through seaports and the international airports from endemic countriesinto Cameroon without thorough inspection cannot be rolled-out. Every year, transhumantanimals come to the East and Adamawa, from Chad and the Central African Republic.Similarly, every year, several heads of ruminant livestock come on transhumance from neighbouring Chad, Niger and Nigeriato the Farnorth Region of Cameroon thereby increasing animal density and contacts that permit FMD introduction and distribution. Further, these animals entering the country are generally not diagnosed or thoroughly inspected for FMD at the different veterinary checkpoints before allowing them to enter the country. To avoid the introduction and spread of FMD viruses in the national territory, a competent epidemiological surveillance infrastructure is required at different livestock entry points of the country.

Abattoirs and livestock markets: The modern abattoirs located in some Cameroonian cities such as Yaounde, Douala and Ngaoundere belong to the state-owned livestock production and exploitation corporation (SODEPA). However, there are several slaughter slabs with low veterinary service inspection. Information from available sources indicates that the abattoirs and livestock markets in Cameroon have no diagnostic facilities and limited technical capacity to identify some diseases like FMD. However, the veterinary service inspection is based on visual examination. Although the thorough inspection is supposed to be conducted in abattoir and livestock markets, it is rather unfortunate that such services do not exist for FMD as veterinary chief of centres in some major cattle production areas do not have data on FMD cases. Abattoirs have been identified as risk areas for the spread of infectious diseases because it receives animals from everywhere and some of them have been reported to be sick (Mebanga et al., 2020). The handling of animal carcasses requires high biosecurity measures to avoid personal and environmental contamination. If infected animal carcasses, secretions and excretions get into the environment, it could lead to the spread of highly contagious diseases such as FMD in such zones. Samples collected in abattoirs and cattle markets in six locations within the main agro-ecological zones of Cameroon revealed highcontamination in Douala and Bertoua. Genetic analysis of one of the samplesled to the isolation of serotype O of the East African-3 lineage.

Regarding the livestock trading system of Cameroon, it is characterized by trade animals from neighbouring Chad passing through theFar-North Region of Cameroon tothe Borno state in Nigeria. During transit in Cameroon, they mix with resident herds during transhumance towards the Waza park. There, they come in contact with wild reservoirs through the sharing of pasture, drinking and camping areas. The animals coming in through the North and Adamawa regions from the Central African Republic could be contaminated upon contact with those found in the Benoue and Bouba-Njidda game reserves on their way to the Adamawa state of Nigeria. Besides transit to Nigeria, there is transit to Gabon and Equatorial Guinea across all intervening regions to the South region. Major cattle markets of Cameroon receives animals from different regions and neighbouring countries with some of the animals subclinically and clinically infected. The animals are transported to markets on foot and by lorries. Animals from different herds are often packed in the same lorry. Similarly, during trekking to livestock markets, two or more animal herds usually come in contact either on their way to the market or upon return. During market days a subset of cattle merchants buy animals and they are stocked in one large pen before their transportation from one local market to the next. This kind of market system promotes the spread of infectious diseases especially FMD between localities. It should be noted that not all animals taken to these markets are sold and if they come in contact with sick counterparts and are carried home, they could contaminate other susceptible ones.

Fomites: The implication of fomites in FMD spread is another avenue of research that is neglected globally. The factors suspected to be responsible for FMD spread include various domestic/wild animals, birds, men, truck tires and postal materials. Persistence of the FMDV on surfaces is a potential transmission risk and include animal farms (49 weeks), soil (1 to 21 weeks), manure (1 to 24 weeks), vegetables (1 week), H₂O (3-14), house flies (10 weeks), ticks and haematin of ticks (15-20 weeks), wool of sheep (2 weeks), hair of cattle (4-6 weeks), and cattle skin used as cloth (Sevidzem et al., 2019c). For FMDV to persist for longer periods on these surfaces, optimal environmental conditions must occur. Weather variables such as temperature, relative humidity and pH have been reported to affect FMDV persistence in endemic settings (Mielke and Garabed, 2019).

In 2016 the first author of this paper was interested in knowing the role of stable flies especially the blood-sucking *Stomoxys niger*(Fig. 2a) in the contaminative transmission of FMDV during outbreaks. Preliminary findings of the study conducted during the 2016 FMD outbreak in Ngaoundere of the Adamawa region of Cameroon indicated that this fly vector's mouthparts (Fig. 2b) and legs (Fig. 2c) were contaminated with the FMDV (Fig. 2d) RNA and legs frequently contaminated than mouthparts but mouthparts had a high viral load (Sevidzem et al., 2019c).

The association of the biting predilection sites of different stable flies with areas on the host model where FMD clinical lesions are produced indicated a strong association with these two variables. This shows that flies coming in contact with open sores containing FMDV could lead to their contamination and if they land around open sores of susceptible host this could lead to successful inoculation of the virus (Sevidzem et al., 2020).

Control efforts: The average cost of FMD treatment in Cameroon is estimated at 80, 000 FCFA/herder/year on drugs and the average annual expenditure on drugs for its treatment is estimated at 32,000,000,000 FCFA (32 billion FCFA). There are no vaccination programs and available commercial vaccines to curb the spread of FMD in Cameroon. However, vaccine trials, topical formulation trials and studies to know the different FMD management options by herdsmen have been conducted.



Fig.2. Stomoxys and its body parts involved in the contaminative transmission of FMDV. (a) Adult Stomoxys; (b) Scanning electron microscopy of proboscis; (c) Scanning electron microscopy of tarsus; (d) FMD virus (millions of FMDV particles (size 0.03µm) presumably attach to the hairs of the pulvillus of legs when the fly lands on infected body parts. They could also attach to the peristomal teeth and labellae of probosci when the fly tries to imbibe infected blood, excretions and secretions

Vaccination: The MINEPIA together with LANAVET in 2013, acquired 150 000 doses of an FMD trivalent vaccine prepared by the BVIusing endemic serotypes (A,O,SAT2). The first FMD vaccination campaign in Cameroon was launched in March 2014. This pilot campaign targeted dairy herds in four divisions of two regions-Adamawa (Vina and Mbere divisions) and North-West (Mezam and Bui divisions).The major pitfall of this campaign is that there was no Post Vaccination Monitoring (PVM) record available to say if the vaccine was efficient or not.These weaknesses and challenges in the implementation of FMDvaccination in Africa have already been reported (Maree et al., 2020; Jackson et al., 2021).

Tri-Solfen: TheAustralian wound dressing formulation, TS(Medical Ethics Ptv Ltd, Australia) is registered for use in cattle and small ruminant husbandry in Australia and New Zealand and has recently been registered for FMD therapy in Lao People's Democratic Republic (PDR). This formulation offers numerous advantages over current therapies. TS has been confirmed as providing significant pain relief and more rapid healing of wounds and lesions (Windsor et al., 2020). Further, with a pH of ~2.7, it potentially has a viricidal impact,

avoiding the need for other treatments, including antibiotics. The dressing contains two local anaesthetics (lignocaine and bupivacaine), adrenalin and an antiseptic (cetrimide) in a gel formulation that creates a barrier effect, numbing the pain of lesions, rapidly reducing their infectivity, and hastening to heal, potentially reducing the weight loss in affected individuals.

From the results of the first clinical trial of TS in Cameroon,farmer observations indicated that the TStreatment group had a more rapid return to eating with cessation of excessive salivation, and more rapid return of mobility (walking) with the absence of overt lameness compared to the commonly used antibiotic Moore-Oxy® (MO) (oxytetracycline HCL 5%, Hebei Kexing Pharmaceutical Co. Ltd., Shijiazhuang City, China) and the non-treated control group (Lendzele et al., 2020).100% of farmers expressed their desire that the product is made available for use in their region and cost-effective modelling indicated that TS therapy only costs 2.5 USD, imposes no additionalfinancial burden on farmers, with the treatment likely to be provided at a similar or reduced cost to current treatment choices. Further, TS can have a significant impact on improving lesion treatment and reducing viral transmission during outbreaks, whilst addressing risks of antimicrobial food safety and antimicrobial resistance (AMR) issues.

Farmers own control methods

Chemotherapy: As no FMD vaccination programs, commercial vaccines and TS are available to farmers in Cameroon, they have devised other strategies to manage the disease. A field study in major cattle markets where most drug dealers come to sell their products was surveyed and it was noticed that farmers purchasedantibioticsand antiparasiticsto manage FMD. Only a few of them practised herd hygiene (cleaning of wounds) and herd isolation. Procaine (antibiotic) was the most frequently used management option (51.23%) followed by procaine plus medicinal plants (13.22%) and medicinal plants (10.74%) (Sevidzem et al., 2019b). The most frequently prescribed drugs for the management of FMD were antibiotics followed by antiparasitic and lastly by antiinflammatories. The majority of the drug sellers prescribed antibiotics precisely MO whose common name was boldly written on its bottle as "Likki Bauru" in Fulfude, meaning FMD drug (Sevidzem et al., 2019b).

Husbandry management practice: Another herd management practice that facilitates FMD transmission in herds is the use of sick cases to speedherd transmission so that they can treat them at once to reduce management costs. In the endemic Far-North region of Cameroon with high FMD serotype diversity, the issue of how much cross-protection between FMD serotypes might prevent transmission was investigated for serotypes O and A. From the simulation runs, it was concluded that cross-immunity would not prevent FMD infections in the Cameroonian context (Xiao et al., 2015).

Medicinal plants: Despite the frequent use of different veterinary pharmaceuticals to manage FMD in Cameroon, traditional medicinal plants are also widely used. The treatment of animals with these medicinal plants is often associated with antibiotics (Sevidzem et al., 2019b). In the Far North region of Cameroon12 medicinal plants are commonly used by farmers to manage FMD (Vougat et al., 2015). Interestingly, two of the medicinal plants known as Bosica senegalensis and Tapinanthus dodoneifolius possesses active compounds that are responsible for several biological activities, reason why they are used by farmers to manage FMD. However, only 10% of farmers interviewed in the different cattle markets of Adamawa confirmed the use of only medicinal plants to treat FMD. Although these medicinal plants are widely available, farmers still prefer antibiotics like 'procaine' that is cheap and available than traditional preparations that are difficult to prepare, time-consuming and more expensive.

FMD progressive control pathway for Cameroon: The PCP-FMD is a stepwise framework that assistscountries to progress towards achieving freedom from FMD (with or without vaccination) after official approval by the OIE (Fig. 5). Cameroon is presently in stage one (Ehizibolo et al., 2020) of the PCP-FMD and a risk-based control plan is ongoing. The Cameroonian Government is willing to control FMD and this is manifested through the training of staff of Divisional delegations. The Government also offersfinancial support to LANAVET to boost its testing capacity. To improve its diagnostic infrastructure to satisfy the increasing demands, LANAVET is currently creating branches in the different regions of the country. There are no records of FMD regional control initiatives in Central Africa, but the Government's commitment to other regional control initiatives against epizootics is evidence of its readiness in case of an eventual regional control initiative.



Fig. 5:The different PCP stages for FMD control with Cameroon in stage 1.

Gaps analysis

*Epidemiological surveillance studies:*Apart from the comprehensive national study on FMD carried out in 2012, no systematic nationwide data on this disease is available. There is a need to increase the FMD testing capacity, to conduct regular field surveys at regional and border levels togenerate robust epidemiological data.

Socio-economic impact studies: There is a need to update the socio-economic impact of FMD on the different livestock breeder categories.

Risk-based studies: There is a need to study the role of wild animals in FMD dissemination as studies on this subject has not yet been realised in Cameroon. The role of small ruminants in FMD transmission in mixed herds with cattle should also be studied. The role of environmental contaminants such as soils, air at an abattoir, cattle markets, and herds as well as other fomites (swabs from lorries transporting cattle from one cattle market to the next, livestock owners and health workers) should be conducted to design biosecurity measures typical of the Cameroonian setting, required to curb the spread of FMD. As preliminary studies have already indicated that stable flies could serve as mechanical vectors, there is a need to collect samples during an outbreak and non-outbreak periods to determine virus

Source: Modified from https://www.fao.org/ag/AGAinfo/commissions/docs/P CP/PCP en.pdf.

Abbreviations: FMD, foot-and-mouth disease; OIE, World Organisation for Animal Health; PCP, Progressive Control Pathway.

persistence and to conduct further experiments to define their vectorial capacity.

FMD control studies: The FMD control phase in Cameroon will require that robust, credible and updated national epidemiological data is available to guide tailored control. For this, researchers, state veterinarians and those in the private sector could contribute to generating such information. The lessons learned from the last vaccine campaign could help inform future vaccine trial efforts in Cameroon. The first field trial of TS was very successful and this product is already authorised for use in Cameroon (authorisation number: MA nº DRL0101120/CMR).

Conclusion

Four FMD serological types (SAT 1, SAT 2, O, A) circulates in the animal populations of Cameroon. Abattoir and cattle markets environments represent transmission risk areas. Stomoxys niger could be involved in FMD spread. Questionnaire studies revealed the possibility of animal contact during transhumance and the difficulty faced by farmers to quarantine their animals during outbreaks to curb the spread of FMD. No vaccination programs and commercial vaccines are available in Cameroon. The first tri-solfen clinical trial was very successful and was authorised for use in Cameroon. The main gaps in FMD research are lack of regular surveillance data, no comprehensive socio-economic FMD impact studies, studies on the role of wild animal hosts in its transmission is not available.

Reference

- Abdel-Aziz AI, Aurore R and Relmy A (2019). Seroprevalence and molecular characterization of footand-mouth disease virus in Chad. Vet. Med. Sci.00 : 1–8.
- Atuman, YJ, Kudi CA and Abdu PA (2020). Seroprevalence of Foot and Mouth Disease Virus Infection in Some Wildlife and Cattle in Bauchi State, Nigeria. Vet. Med. Int. 8.
- Bertram MR, Bravo de Rueda C and Garabed R (2018b). Molecular epidemiology of foot and mouth disease virus in context of transboundary animal movement in far North Region of Cameroon. Front. Vet. Sci., 5 : 320.
- Bertram MR, Delgado A and Pauszek SJ (2018a). Effect of vaccination on cattle sub-clinically infected with foot-and-mouth disease virus

in Cameroon. Prevent. Vet. Med., 155 : 1-10.

- Bertram MR, Dickmu S and Palinski RM (2019). Genome sequences of four foot-and-mouth disease virus SAT 1 topotype X isolates from Cameroon. Microbiol. Resour. Announc. 8 : e01243-19.
- Bronsvoort BM, Handel IG and Nfon CK (2016). Redefining the "carrier" state for foot-andmouth disease from the dynamics of virus persistence in endemically affected cattle populations. Sci. Rep. 6 : 29059.
- Bronsvoort BM, Anderson J and Corteyn A (2006). Geographical and Age-Stratified distribution of Foot and Mouth disease virus-seropositive and probang-positive cattle herds in the Adamawa province of Cameroon. Vet. Rec. 158 : 299-308.
- Bronsvoort M, Tanya VN and Kitching RP (2003). Foot and mouth disease and livestock husbandry practices in the Adamawa Province of Cameroon. Trop. Anim. Health. Prod., 35 : 491–507.
- Bronsvoort MC, Nfon C and Hamman SM (2004). Risk factors for herdsman-reported footand-mouth disease in the Adamawa Province of Cameroon. Prevent. Vet. Med., 66 : 127–39.
- Buckle K, Bueno R and McFadden A (2021). Detection of Foot-and-Mouth Disease Virus in the Absence of Clinical Disease in Cattle and Buffalo in South East Asia. Front. Vet. Sci., 8 : 691308.
- Chepkwony EC, Gitao GC and Muchemi GM (2021). Epidemiological study on foot-andmouth disease in small ruminants: Seroprevalence and risk factor assessment in Kenya. PLoSONE. 16 : e0234286.
- Colenutt, C (2020). FMD surveillance at markets and abattoirs in Cameroon. European Commission for the control of FMD. EuFMD virtual meeting 27 november 2020.
- Di Nardo A, Libeau G and Chardonnet B (2015). Serological profile of foot-and-mouth disease in wildlife populations of West and Central Africa with special reference to *Syncerus caffer* subspecies. Vet. Res., 46 : 77.
- Dickmu S, Nsangou S, Kouamo J, Poueme R, Awah NJ, Garabed R, Abdoulkadiri S, Rodriguez L and Arzt J (2019). Foot-and- Mouth Disease in Domestic Ruminants in the North and Far North Regions of Cameroon. Global Footand-Mouth Disease Research Alliance 2019

Scientific meeting in Bangkok, Thailand 25th -31st October.

- Ehizibolo DO, Fish IH and Brito B (2020). Characterization of transboundary footand-mouth disease viruses in Nigeria and Cameroon during 2016. Transbound.Emerg. Dis. 00 : 1–14.
- Ekue N, Tanya V and Ndi C (1990). Foot-andmouth disease in Cameroon. Trop. Anim. Health Prod., 22 : 34–36.
- Ekwem D, Morrison TA and Reeve R (2021). Livestock movement informs the risk of disease spread in traditional production systems in East Africa. Sci. Rep., 11 : 16375.
- FAO (2015). Strategic plan for the control of footand-mouth disease in Cameroon. FAO. 67p.
- Jackson B, Harvey Y and Perez-Martin E (2021) .The selection of naturally stable candidate foot-and-mouth disease virus vaccine strains for East Africa. Vaccine., 39 : 5015– 5024.
- James AD and Rushton J (2002). The economics of foot-and-mouth disease. Rev. Sci.Tech., 21: 637-644.
- Jori F and Etter E (2016). Transmission of foot and mouth disease at the wildlife/livestock interface of the Kruger National Park, South Africa: Can the risk be mitigated? *Prevent.* Vet. Med., 126 : 19–29.
- Kerfua SD, Shirima G and Kusiluka L (2018). Spatial and temporal distribution of footand-mouth disease in four districts situated along the Uganda-Tanzania border: Implications for cross-border efforts in disease control'. Onderstepoort J. Vet. Res., 85 : a1528.
- Kim H, Xiao N and Moritz M (2016). Simulating the Transmission of Foot-And-Mouth Disease Among Mobile Herds in the Far North Region, Cameroon. J. Artif. Soc. Soc. Simul. 19.
- Knowles NJ and Samuel AR (2003). Molecular epidemiology of foot-and-mouth disease virus. Virus Res., 91 : 65–80.
- Lendzele SS, Armel KA and Rodrigue MN (2021). Clinical Foot-and-Mouth Disease in Non-Vaccinated Smallholder Dairy Cattle in Adamawa Region, Cameroon: Prevalence, Farmer's Knowledge and Practices. J. Vet. Med. Surg., 5:4-40.
- Lendzele SS, Mavoungou JF and Kong AB (2020).

Efficacy and application of a novel topical anaesthetic wound formulation for treating cattle with Foot-and-Mouth disease: a field trial in Cameroon. Transbound. Emerg. Dis., 00 : 1–12.

- Ludi A, Ahmed Z and Pomeroy LW (2016). Serotype Diversity of Foot-and-Mouth-Disease Virus in Livestock without History of Vaccination in the Far North Region of Cameroon. Transbound. Emerg. Dis., 63: e27-e38.
- Maree FF, Kasanga CJ and Scott KA (2020). Challenges and prospects for the control of foot-and-mouth disease: an African perspective. Vet. Med. : Res. Rep., 202 : 168-194.
- Mebanga AS, Beloko SL and Mamoudou A (2020). Prevalence and risk factors of foot-andmouth disease in the cattle market of Garoua-Boulai and in the transhumance zone of Lome and Djerem in East Cameroon. Int. J. Biol. Chem. Sci., 14: 2799–2806.
- Mielke SR and Garabed R (2019). Environmental Persistence of Foot and Mouth Disease Virus Applied to Endemic Regions. Transbound. Emerg. Dis., 1–12.
- Morgan KL, Handel IG and Tanya VN (2014). Accuracy of herdsmen reporting versus serologic testing for estimating foot-andmouth disease prevalence. Emerg. Infect. Dis., 20: 2048–54.
- Motta P, Porphyre T and Handel I (2017). Implications of the cattle trade network in Cameroon for regional disease prevention and control. Sci. Rep., 7: 43932.
- Motta P, Thibaud P and Saidou MH (2018). Cattle transhumance and agropastoral nomadic herding practices in Central Cameroon. BMC Vet. Res., 14 : 214.
- Moumini (2011). Evolution et situation actuelle de l'élevage au Cameroun, Rapport de synthèse, MINEPIA – Cameroun. 3p.
- Nyaguthii DM, Armson B and Kitala PM (2019). Knowledge and risk factors for foot-andmouth disease among small-scale dairy farmers in an endemic setting. Vet. Res., 50 : 33.
- Paton DJ, Gubbins S and King DP (2018). Understanding the transmission of foot-andmouth disease virus at different scales. Curr. Opinion Virol., 28 : 85–91.
- Pomeroy LW, Bjørnstad ON and Kim H (2015).

Serotype-Specific Transmission and Waning Immunity of Endemic Foot-and-Mouth Disease Virus in Cameroon. PLoSONE. 10 : e0136642.

- Profitós JMH, Moritz M and Garabed RB (2013). What to do with chronically sick animals? Pastoralists' management strategies in the far north region of Cameroon. Pastoralism. 3:8.
- Sansamur C, Arjkumpa O, Charoenpanyanet A and Punyapornwithaya V (2020). Determination of Risk Factors Associated with Foot and Mouth Disease Outbreaks in Dairy Farms in Chiang Mai Province, Northern Thailand. Animals. 10 : 512.
- Schnell PM, Shao Y and Pomeroy LW (2019). Modeling the role of carrier and mobile herds on foot-and-mouth disease virus endemicity in the Far North Region of Cameroon. Epidemics., 29 : 100355.
- Sevidzem SL, Mamoudou A and Dickmu S (2019c). Risk Factors for the Contamination of Wild *Stomoxys niger niger* Macquart 1851 (Diptera: Muscidae) with the Foot-and-Mouth Disease Virus. Curr. Res. Agric. Sci., 6:95–108.
- Sevidzem SL, Mamoudou A and Mavoungou JF (2019a). Serological Epidemiology of Footand-Mouth Disease among Sedentary Mixed-species Herds in Adamawa Region, Cameroon. J. Adv. Microbiol., 17 : 2456–7116.
- Sevidzem SL, Mavoungou JF and Mintsa NR (2019b). Veterinary pharmaceuticals sold in cattle markets for the management of footand-mouth disease and flies in Vina division (Adamawa-Cameroon). Dairy Vet. Sci. J., 10 : 2.
- Sevidzem SL, Mavoungou JF, Zinga-Koumba RC and M'batchi B (2020). Alighting Dipterous Insects on Cattle are Associated to

Contaminative Transmission of Foot-and-Mouth Disease During Epidemics in Ngaoundere Cameroon. Entomol. Faunist – Faunist. Entomol. 73 : 132–144.

- Sinkala Y, Simuunza M and Pfeiffer DU (2014). Challenges and Economic Implications in the Control of Foot and Mouth Disease in Sub-Saharan Africa: Lessons from the Zambian Experience. Vet. Med. Int., 12.
- Ularamu HG, Ibu JO and Abenga JN (2020). Incursion of Foot-and-Mouth Disease (FMD) Serotype O East Africa Topotype -3 (O/EA-3) in Nigeria. Nig. Vet. J., 41.
- Vougat Ngom RRB, Foyet HS and Garabed R (2020). Human Health Risks Related to Penicillin G. and Oxytetracycline Residues Intake Through Beef Consumption and Consumer Knowledge About Drug Residues in Maroua, Far North of Cameroon. Front. Vet. Sci., 7 : 478.
- Vougat RRBN, Foyet HS, Garabed RB and Ziebe R (2015). Antioxidant activity and phytochemical constituent of two plants used to manage foot-and-mouth disease in the Far North Region of Cameroon. J. Intercult. Ethnopharmacol., 4 : 40-46.
- Wekesa SN, Sangula AK and Belsham GJ (2015). Characterisation of recent foot-and-mouth disease viruses from African buffalo (*Syncerus caffer*) and cattle in Kenya is consistent with independent virus populations. BMC Vet Res., 11 : 17.
- Windsor PA, Earp F and MacPhillamy I (2020). A new topical therapy for Foot-and-mouth disease addresses animal welfare and other issues. Vet Med : Res Rep., 11 : 99-107.
- Xiao N, Cai S, Moritz M, Garabed R and Pomeroy LW (2015). Spatial and temporal characteristics of pastoral mobility in the far north region, Cameroon: data analysis and modeling. PLoS ONE. 10 :131–9.
